

Language Status in Children with Birth Asphyxia: A Follow Up Study (Brain and Language)

Shanthini, S. V.,

(Audiologist and Speech Language Pathologist): Dept. of Speech, Language and Hearing Sciences
Sri Ramachandra University (SRU)
Chennai 600116, India
sharonviolashanthini@yahoo.com

Abstract—Development of the brain is crucial to facilitate language acquisition. Any form of injury and malfunctioning of the brain such as “Birth Asphyxia” can inhibit language acquisition in children (Jansson-Verkasalo, 2004) and causes delay in speech and language development (D’Souza et al 1981). This follow up study (2011) is an extension of a pilot study conducted by Shanthini et al. (2009) that focused on analysing the language status of children with birth asphyxia without any other associated problems to their individual APGAR score, time at which APGAR score reached 8 and cognition pertaining to language.

Keywords: Birth Asphyxia; Language status; APGAR score; Time at which APGAR score reached 8; Cognition pertaining to Language.

I. INTRODUCTION

Language is the mode by which human communicates ones thoughts or intentions to an individual or address to a group of people (Owens, 2001). Language acquisition is the process whereby one acquires the knowledge of language, which according to psycholinguists comprises finite set of rules (Chomsky, 1965 as cited in Shulman & Capone, 2010). In this process the child is considered as an active agent who needs to unfold the language both in terms of its structures and pragmatic functions. Almost every child language researchers have established that all normal children acquire language to which they are exposed to, and conversely children who are not normal in one sense or the other find it hard to acquire language (Wiig & Secord, 1982; Mogford & Bishop, 1988). They have identified several contributing factors towards language acquisition such as presence of functionally normal cortical structures, normal cognitive functioning, normal sensory input system, normal motor output mechanism, adequate linguistic environment and healthy child during prenatal, perinatal and postnatal period. Language acquisition is affected when the pre-requisites for normal speech and language development are affected or inadequate.

Development of the brain has been considered as one of the crowning achievements in human evolution which facilitates language acquisition. Any form of injury and malfunctioning of the brain can inhibit language acquisition in children and in case of adult can lead to dysfunction of language (Lenneberg, 1967 as cited in Lust, 2006). Since the language representation and organization is a cerebral process, pathological conditions in it would lead to severe cognitive dysfunction. However, the cognitive deficit in an adult would have quite different consequences when compared to children (Anderson, Smith, Leventer, Coleman, Anderson, Williams, Greenham & Jacobs, 2009). In cases of focal left hemisphere


insult or complete removal of left hemisphere, children show a better recovery pattern than adults (Heywood & Canavan, 1987; Taylor & Alden, 1997). In contrast, children sustaining generalized cerebral insult such as ‘Traumatic Brain Injury’ (TBI) or early ‘Cerebral Vascular Accident’ (CVA) display slower recovery and poorer outcome in intellectual and academic achievements than adults with similar insults (Taylor, Yeates, Wade, Drotar, Stancin & Minich, 2002; Anderson, Catroppa, Morse, Haritou & Rosenfeld, 2005; Ballantyne, Spilkin, Hesselink & Trauner, 2008). It is seen that brain insult prior to the age of 2 years leads to poor cognitive outcome but when the insult occurs at the later childhood, there will have lesser impact in cognition (Anderson et al., 2005). Hence it is evident that normal functioning of the brain is critical for language acquisition.

Generally it is observed that high risk infants are vulnerable to delay in language acquisition due to injury or malfunctioning of the brain (Jansson-Verkasalo, 2004). Birth asphyxia is one of the major high risk factor which has been reported to cause brain insult at the time of birth (Majeed, Menon, Shaikh, Majeed & Rajar, 2007; Haider & Bhutta, 2006); (Windle, 1969) which causes delay in speech and language development (D’Souza, Nolan, McCartney & Taylor, 1981).

Birth asphyxia, also called Asphyxia Neonatorum, is defined as the failure to start regular respiration within a minute of birth. It is a condition in which extreme decrease in the concentration of oxygen in the body accompanied by an increase in the concentration of carbon dioxide that leads to loss of consciousness or death (Cheung & Robertson, 2000; Donn, Sinha & Chiswich, 2002). When this condition is prolonged or the impact of the condition is adverse it leads to injury or malfunctioning of the brain. This will inhibit the aerobic metabolism in the internal structures where the oxygen supply is reduced (Park, Sanders & Maltepe, 2010).

Figure 1 below shows the pathophysiology of birth asphyxia. When this condition occurs it will trigger any one of the two reactions namely ionic shift and depolarization or energy failure and acidosis simultaneously. Once any one of the reaction takes place it will lead the other two reactions interchangeably namely increase in the release of neurotransmitter and there will be a reduction in the uptake of neurotransmitter simultaneously and phospholipid degradation followed by increase in the free fatty acids and free radical lipid peroxidation. All these change in the cellular mechanism results in disrupted cell membranes and metabolic machinery.

Figure 1



This is why children who had birth asphyxia for a prolonged period is most likely to exhibit neurological impairment, motor impairment, cognitive impairment, language deficits, speech and language disorders, sensory impairments or other consequences, manifestations or associated problems depending on the site of oxygen deprivation.

The incidence of birth asphyxia varies across different countries, states, urban and rural areas. This also depends on biological factors such as antenatal care which include prenatal (poor nutritional status and anaemia), perinatal, post-natal care and others (Majeed et al, 2007). Even environmental factor such as socio- economic status plays a vital role in the incidence of birth asphyxia (Straube, Voigt, Jorch, Hailler, Briese, & Borchardt, 2010). The incidence of asphyxia in Britain in 1960's and in the United States in 1970's was 1.2 to 5% of birth asphyxia among which 0.4% to 1.6% had severe form of asphyxia (Addy, 1982). Of 25 million infants born in India, the incidence of asphyxia was found to be 3 to 5% (Ghai, Gupta, & Paul, 2004).

There are many causes of birth asphyxia. The most common cause are prenatal hypoxia, fetal distress, umbilical cord compression, knotting of umbilical cord around the head, neck or body of the faetus, aspiration of amniotic fluid, meconium stained and others (Majeed et al, 2007; Simon & Morley, 2005).

The clinical signs of birth asphyxia are based upon the colour, heart rate, reflex irritability, muscle tone and respiratory effort. The symptoms of asphyxia are cyanosis, bradycardia, hypotonia, poor response to stimulation and reduced respiratory effort.

Diagnosis of birth asphyxia can be objectively made by using the APGAR score, a recording of physical health of the newborn, determined after the examination of the appearance, pulse rate, grimace, activity and respiration (Apgar, 1953). APGAR score is obtained at the 1st, 5th, 10th, 15th and 20th minute of birth. However the APGAR score of the 1st, 5th and 10th minute is generally taken to check whether there is any neurological impairment. Occasionally children with birth asphyxia gets an APGAR score of 8 at the 2nd minute of birth itself and in those cases a mention of that time will be noted in their discharge summary (Khreisat & Habahbeh, 2005).

Generally the scoring pattern of calculating the APGAR score depends on the symptoms and associated problems of each signs. The independent score is given based on the severity of the symptoms and associated problems manifested for each signs exhibited by the baby at the point of birth. The APGAR scoring system chart is illustrated in the table 1.

Table 1

APGAR five components	SIGNS	SYMPTOMS AND ASSOCIATED PROBLEMS			TOTAL SCORE ACROSS TIME IN MINUTES				
		Score =0	Score =1	Score =2	1	5	10	15	20
Appearance	Colour	Blue or Pale	Acro-cyanotic	Totally Pink					
Pulse	Heart Rate	Absent	<100 Beats/Minute	>100 Beats/Minute					
Grimace	Reflex Irritability	No Response	Grimace	Cry or active Withdrawal					
Activity	Muscle Tone	Limp	Some flexion	Active motion					
Respiration	Respiratory Effort	Absent	Weak Cry	Good crying					

APGAR score of 8 at the 2nd min of birth: (If applicable)

The APGAR scoring system chart

Table 2 below indicates the scoring pattern of APGAR score which is done by adding the corresponding score of each sign in the APGAR score sheet. Since there are 5 clinical signs, the total score will be out of 10.

Table 2

Symptoms and associated problems	Score
Normal symptoms	2
Slightly deviant	1
Deviant	0

The scoring pattern in calculating the APGAR score

The severity of birth asphyxia is obtained by the APGAR score as shown in Table 3.

Table 3

APGAR score	Severity
8 to 10	Normal
5 to 7	Mild
3 to 4	Moderate
0 to 2	Severe

The severity scale of birth asphyxia

Birth asphyxia can cause various consequences depending on where is the extent of oxygen deprivation, exact site of lesion, APGAR score at the time of birth and severity of birth asphyxia. The consequences of birth asphyxia are language impairment such as “Delayed Speech and Language Development” (D’Souza et al, 1981) and “Specific Language Impairment” (Stanton-Chapman, Chapman, Bainbridge & Scott, 2002), neurological impairment such as “Learning Disability” (Bhate & Wilkinson, 2006), “Autism Spectrum Disorders” (Simon et al, 2005), “Attention Deficit Hyperactive Disorder” (Gustafsson & Kallen, 2010) and “Seizural Disorder” (Airede, 1991), neuromotor speech disorder” such as “Cerebral Palsy” (Simon et al, 2005), fluency disorder such as “Stuttering” (Somefuna, Lesib, Danfulania & Olusanyac, 2005), sensory impairment such as “Hearing Impairment” (Ohl, Czajka, Chobaut & Tavernier, 2009) and “Visual Impairment” (Ronald, Jan, Hill & Wong, 1986), cognitive impairment such as “Mental Retardation” (Simon et al, 2005), mortality (Jerneck & Herbst, 2001) and other complication either after birth or in the later stages of their lifetime. These manifestations can occur in isolation or in combination. Children with birth asphyxia are prone to have associated problems (Avery & Taeusch, 1984). This high risk factor, “Birth asphyxia” can occur in isolation or in combination with

other associated high risk factors such as low birth weight, preterm, hyperbilirubinemia, and others.

II. REVIEW OF LITERATURE

Research relating Birth Asphyxia to Neonatal Encephalopathy, Motor and Cognitive Impairment

Gonzalez and Miller (2006) found that children who survived from neonatal encephalopathy following perinatal asphyxia exhibited cognitive impairment. However for those who have survived from severe neonatal encephalopathy exhibited both motor and cognitive impairment.

Research relating Birth Asphyxia to Low APGAR Score, and Neurological Impairment (Cerebral Palsy) and Associated High Risk Factor (Low Birth Weight)

Paneth (2010) stated that children who had normal weight and had a low APGAR score of 4 for a prolonged period of time had cerebral palsy. However in children who had low APGAR score and with an associated problem of low birth weight, cerebral palsy was less predictable because the presence or absence of cerebral palsy cannot be attributed directly to the low APGAR score alone. However the time at which the APGAR score reached 8 was brief or prolonged, APGAR score had improved, reduced or remain constant with respect to time and the severity of the other associated problem should be taken into account.

Research relating Birth Asphyxia to Cognitive and Behavioural Impairment

Anderson et al. (2009) found that children who had early brain insult before age 2 years exhibited global and significant cognitive deficits whereas children who had early brain insult after age 2 functioned closer to normal expectations. A linear correlation was noticed between the age of insult and the outcome. On the other hand when the behavioural domains were assessed, it was seen that children with late early brain insult from 7 to 9 years performed worse than those with early brain insult from 3 to 6 years. This concluded that not all functions share the same pattern of vulnerability with respect to age at insult.

Research relating Birth Asphyxia to Neonatal Encephalopathy, Cognitive Impairment, Academic Performance, Neuropsychological Functioning and Behavioural Impairment

Handel, Swaab, Vries and Jongmans (2007) found that children with relatively mild degree of encephalopathy due to birth asphyxia performed relatively better in the 4 domains tested (cognition, academic, neuropsychological functioning and behaviour) when compared to those with severe encephalopathy. However those who had moderate neonatal encephalopathy had varying outcomes.

Research revealing Language Status in children with High Risk Factors

Chelvi (2007) found that the combined language age of the children with high risk such as very low birth weight and low birth weight with preterm children was delayed by 9 months when compared to the typically developing children.

Research relating Birth Asphyxia to Neonatal Encephalopathy, Low APGAR score before and after 5th Minute of Birth, Cognitive Impairment and Educational Performance

Odd, Rasmussen, Gunnell, Lewis and Whitelaw, (2008) found that children who had brief low APGAR scores before the 5th

minute of birth and those who had prolonged low APGAR scores after 5th minute of birth had low IQ scores at the age of 18 years which is mainly exhibited at the later period of life as the educational demand increases.

Research relating Birth Asphyxia to Neurological Impairment, Psychological Status and Quality of survival

Thomson, Searle and Russell (1977) investigated the neurological and psychological status of the 31 children who had survived from severe birth asphyxia and was compared to their age matched control group. Out of 31 children, 29 did not exhibit any serious neurological or mental handicap where as 2 of the children had severe neurological and cognitive impairment. Hence quality of life will be improved in children who had survived from severe birth asphyxia when resuscitation is encouraged.

Research relating Birth Asphyxia to 5th Minute APGAR Score, Socio Economic Status and Biological Factors

Straube et al. (2009) investigated the association of 5th minute APGAR score with maternal socio-economic and biological factors and found that, a low APGAR score was commonly noticed in overweight women and in women who were above 35 years of age where as socio-economic factors did not significantly influence APGAR scores.

Research relating Birth Asphyxia to Low APGAR Score and Maternal Socio Economic Status

Odd, Doyle, Gunnell, Lewis, Whitelaw and Rasmussen (2008) found that children who had the risk of a low APGAR score were more in the mothers who had low educational status and lack of awareness on birth asphyxia as a resultant of poor socio economic status.

Research relating Birth Asphyxia and Preterm to Cognitive Impairment and Language Impairment

Golightly, Sander and Raz (2003) revealed that children who had more than one high risk factors will exhibit have severe cognitive and language impairment when compared to those with a single high risk factor.

Research relating Risk Factors Birth Asphyxia and Early Identification

Majeed et al. (2007) found the risk factors of birth asphyxia in neonates were increased or decreased maternal age, poor antenatal care, multiple births, anaemia, poor nutritional status, other intrapartum risk like meconium stained and other complications. It was found that the mortality rate reduced when there was early identification of high risk cases and improved antenatal and perinatal care.

Research relating Low APGAR Score to Mortality, Neurological Impairment and Cognitive Impairment

Jerneck and Herbst (2001) found that infants with very low APGAR score at the 5th minute exhibited mortality, cognitive impairment or neurological impairment. The incidence rate varied depending on environmental factors.

Research relating Very Low 5th Minute APGAR Score and Preterm to Mortality

Lee, Subeh and Gould, (2010) found that neonates who had very low 5th minute APGAR score (0 to 3) and were born at the 24th to the 28th week of gestation had increased mortality rate when compared to the those who were born after 28th week of gestation.

Research relating Very Low 1st Minute APGAR Score, Preterm and Very Low Birth Weight to Mortality

Genzel-Boroviczeny, Hempelman, Zoppellian and Martinez (2010) found that despite frequent intubations, preterm children who were born within 23 to 26 weeks of gestation with very low birth weight (500g) and very low APGAR score (0 to 1) at the 1st minute of birth exhibited higher mortality rate especially males than those with an APGAR score above 1 at the 1st minute of birth.

Research relating Birth Asphyxia to Neonatal Encephalopathy, Cognitive Impairment and Educational Performance

Odd, Whitelaw, Gunnell and Lewis (2010) found that among the children who survived neonatal encephalopathy due to birth asphyxia, the children with symptomatic neonatal encephalopathy after resuscitation had exhibited cognitive impairment, poor language skills and poor educational attainment.

Research relating birth asphyxia and the risk factors of neonatal mortality

Lee, Mullany, Tielsch, Katz, Khatri, LeClerq and Darmstadt (2008) found that 9.7 out of 1000 live births died in southern Nepal in the year 2002 to 2006 due to birth asphyxia, maternal infections, premature births, multiple birth and poor socio-economic status of the parents.

Research relating birth asphyxia and hearing loss

Ohl et al. (2009) found that the sole cause of sensorineural hearing loss was attributed to severe birth asphyxia, neurological disorder, syndromes associated to cause of hearing loss, TORCH infections, any family history of deafness or when the infants have more than one high risk factor at birth such as low birth weight and preterm. However conductive hearing loss was found in infants with craniofacial anomalies such as cleft palate and ear aplasia.

Research relating Birth Asphyxia and Learning Disability

Bhate et al. (2006) found that the causes of learning disability can be genetic such as down syndrome and fragile-X syndrome, biological factors such as perinatal factors such as birth asphyxia, mechanical birth trauma, hypoglycaemia and prematurity, environmental and idiopathic.

Research relating birth asphyxia and visual impairment

Ronald et al. (1986) used instrumental analysis to detect the presence of visual impairment in children with birth asphyxia since these children are poorly identified due to involvement of multiple neurological impairments. The researchers found that by using cranial computed tomography and visual evoked potential mapping, children with cortical visual impairment were identified. The researcher could associate the effects of birth asphyxia to cortical visual impairment.

Research relating Birth Asphyxia and Attention Deficit Hyperactive Disorder

Gustafsson et al. (2010) found that children who were born in Sweden and diagnosed with Attention Deficit Hyperactive Disorder had biological high risk factors such as birth asphyxia (APGAR score of less than 7 at the 5th minute of birth) and preterm (less than 32 weeks of gestation) and environmental risk factors such as young maternal age and maternal smoking.

Research relating stuttering and birth asphyxia

Somefun et al. (2005) investigated the causes for children with communication disorders such as hearing impairment, speech disorders (stuttering), language disorders and rhinolalia in the age range of 6 months to 15 years were birth asphyxia, seizures, meningitis, kernicterus, ototoxicity, otitis media effusion, cerebral palsy, measles, congenital deformity and others.

Research relating Birth Asphyxia and Seizural Disorder

Airede (1991) investigated 57 infants with neonatal seizures and found the cause of the neonatal seizures in these children was due to perinatal asphyxia, hypoglycaemia and preterm. However the outcome of the preterm children was poorer than the others.

Research relating Birth Asphyxia to Speech, Language and Hearing Issues

D'Souza et al. (1981) found that among the 26 children who have survived from severe perinatal asphyxia, only 1 child had sensori-neural deafness, whereas one-third of the children exhibited deficits in speech and language without any serious mental or physical handicap. However the authors suggested that the quality of would be better if detected early and treated adequately.

Studies relating Birth Asphyxia and Language Impairment

Stanton-Chapman et al. (2002) stated that children with specific language impairment had risk factors such as very low birth weight and birth asphyxia at birth and the environmental risk factors such as low maternal education, late or no prenatal care and high birth order.

Research relating Multiple Risk Factors and Early Language Development

Sidhu, Malhi and Jerath (2010) found that children who had biological risk factors at birth such as birth asphyxia, preterm, low birth weight, neonatal jaundice and environmental risk factors such as low maternal and paternal education status, low income, higher birth order, large family size, disadvantageous caste, absence of father, low level of occupation of the head of the family exhibited delayed language development.

Research relating Birth Asphyxia and Language

Beharelle, Dick, Josse, Solodkin, Huttenlocker, Levine and Small (2010) found that the children with early unilateral left focal brain injury children had either bilateral activation left and right superior temporal inferior parietal regions or unilateral activation left or right superior temporal inferior parietal regions instead of the left frontal and lateral temporal regions which is essential for language. The outcome was better in those with bilateral activation in terms of processing language.

Research relating the association of Birth Asphyxia to the APGAR Score, Time at which the APGAR Score reach 8, Cognition Pertaining to Language and Language Status

Shanthini and Athmacharan (2009) did a pilot study on the language status in the children with birth asphyxia. In this study, 640 children with birth asphyxia were subjected to a range of evaluations such as neurological evaluation, psychological evaluation, audiological evaluation and detailed speech and language to rule out associated problems. Based on that, 15 children who met the inclusion criteria constituted the subjects of the study who were in the age range of 2 ½ to 4

years. All these children had an APGAR score of 7 at the 1st minute after birth. From the study, it was found that when the APGAR score reached 8 at 2nd minute there was no language delay. When the APGAR score reached 8 at the 5th minute there was a delay of 1 year in the combined language level. When the APGAR score reached 8 at 10th minute there was a language delay of 15 to 24 months.

III. NEED OF THE STUDY

In the last three decades several systematic studies have been carried out in children with birth asphyxia. Most of the researchers have focused on either a single parameter and an issue or a group of parameters and the issues as a resultant of birth asphyxia. Hardly any study has been carried out in documenting the language status in the children with birth asphyxia by giving importance to the role of APGAR score, time at which the APGAR score reached 8, cognition pertaining to language and the language status. Even the little research done in these children has not been followed up to establish the language status of the children as they get older. Also comparative study, if any, between normally developing children and children with birth asphyxia have not been attempted. Besides that, there exists hardly any study that compares the language amongst gender. However in this study cognition is also given equal importance because it precedes language development (Piaget, 1954 as cited in Owens, 2001). So in order to address these issues mentioned above, the following objectives were set.

IV. OBJECTIVES OF THE STUDY

1. To check whether there is any deviance in the language status of the children with birth asphyxia in relation to their APGAR score, time at which the APGAR score reached 8 and their cognitive level pertaining to language as the subject's age increases.
2. To compare the language status of the children with birth asphyxia to that of the typically developing children with respect to their gender.

V. METHODOLOGY

This follow up study was done for dissertation, which was an extension of a pilot study conducted by Shanthini et al. (2009) that focused on analysing the language status of children with birth asphyxia without any other associated problems. Shanthini et al. (2009) reviewed 640 children with birth asphyxia (APGAR score of 7 or lesser at the 1st minute of birth) who were born in a single hospital in Chennai between 2004 and 2005. The records of these children were scrutinized and a series of evaluations such as neurological evaluation (Reflexes, Muscle tone and Motor abilities), psychological evaluation (Geschells Children Behavioural scale and Vineland Social Maturity Scale), audiological evaluation (Visual Reinforcement Audiometry or Conditioning Audiometry and Immittance screening) detailed speech and language (Modified 3 Dimensional Language Acquisition Test). Out of the 640 children who had birth asphyxia, only 15 children (9 males and 6 females) in the age range of 2 ½ to 4 years did not exhibit any associated problems. A detailed speech and language assessment on the 15 children revealed that 10 of them exhibited a delay in language acquisition. The present study is a follow up of all those 15 children after 1 ½ years to check the present language status in generating certain morpho-syntactic structures by comparing them with age and gender matched typically developing children who were also screened with the series of evaluations.

Subject selection: Clinical group

The clinical group for this study included all the 15 children with birth asphyxia (APGAR score of 7 at the 1st minute of birth) without any associated problems identified in the pilot study. All the subjects were Tamil speaking children and did not attend any formal speech and language therapy. All these children were in the age range of 48 to 66 months, with a mean age of 4.38 years and a standard deviation of 0.481.

Table 4

Parameters	Mean	Standard Deviation
Chronological Age	4.38 years	0.481
Developmental Quotient	93	0.755
Social Maturity	4.57 years	0.506

The mean and standard deviation of the Chronological Age, Developmental Quotient & social maturity of the children in the clinical group

The subjects were further classified based on the time at which they reached a normal APGAR score of 8. Of the 15 children, 5 children obtained an APGAR score of 8 at the 2nd minute of birth, another 5 children obtained an APGAR score of 8 at the 5th minute of birth and remaining 5 children obtained an APGAR score of 8 at the 10th minute of birth. All the children have migrated to areas in and around Thiruchirapalli and Madurai districts of Tamilnadu and are presently studying kindergarten and 1st standard (primary 1) under the government sector.

Control group

An equal number of children, matched for age and gender were recruited to form the control group of the study. The mean age of these children was 4.38 years with a standard deviation of 0.481.

Table 5

Parameters	Mean	Standard Deviation
Chronological Age	4.38 years	0.481
Developmental Quotient	93.26	0.703
Social Maturity	4.512 years	0.481

The mean and standard deviation of the Chronological Age, Developmental Quotient & social maturity of the children in the clinical group

All the children in the control group were Tamil speaking children, recruited from a kindergarten and primary school in Chennai. These children were screened to rule out any associated problems such as neurological impairment, sensory impairment, motor impairment, cognitive impairment and delay in speech and language. In total, 30 children participated in this study, 15 each in the clinical and control group.

Tasks given and material development

The deviances of language, if any, in the children with asphyxia were evaluated in relation to certain pre-selected morpho-syntactic structures. For this purpose, the tasks of general conversation and picture & live description were employed. In both the tasks, the use of case markers, tense markers, person marker, number and gender marker was targeted. The picture & live description task included answering targeted questions related to 46 black and white line drawn pictures and 4 live situations. These picture cards were taken from the study of Kavitha (2009), which focused on analysing verb and noun morphology in spoken and written language of children with and without learning disability using Tamil as the native language. Line drawn black and white picture cards were used to avoid distraction in the children. In total, the description task included 50 test items with 50 questions. In the general conversation task, 50 questions were framed to elicit responses that would have the target morpho-

syntactic markers as responses. This was included to verify whether each child produced the same structures in both the tasks or behaved differently in both the tasks. Totally 100 questions were presented to both the clinical group and to the control group.

version 14.0. The score obtained in both the tasks were represented in percentage.

Statistical analysis

Pearson’s correlation was used to investigate the relation between the time at which the APGAR score reached 8 to the performance of the children in the picture description task and general conversation tasks as their age increases. Independent sample t-test was used to investigate the role of gender differences in the performance obtained by the children from both the tasks. Post hoc analyses-Tukey’s HSD (High Significant Difference) test was used to check the language deviance of the children with birth asphyxia by comparing their performance in both the tasks with the age and gender matched control group.

Table 6

Category No.	Category Name	Stimulus No.	Marker's Name	Marker
1.	Case Markers	1 & 2 3 & 4 5 & 6 7 & 8 9 & 10 11 & 12 13 & 14	Accusative Dative Instrumental Sociative Ablative Genitive Locative	-ai -kku -a:l -oda/-kuda -ilirundu -o:da -il
2.	Present Tense Marker	15 & 16	-	-kkir-
3.	Future Tense Markers	17 & 18 19 & 20	-	-pp- -v-
4.	Past Tense Markers	21 & 22 23 & 24 25 & 26 27 & 28 29 & 30	- - - - -	-tt- -nt- -d- -in- -i-
5.	Person/ Pronoun Markers	31 32 33 34 35 36 37 38 & 39 40	1 st person singular 1 st person plural 2 nd person singular 2 nd person plural 3 rd person masculine 3 rd person feminine 3 rd neuter 3 rd person honorific singular 3 rd person honorific plural	na:n na:ngal ni: ni:ngal a:van a:val adu avar/ar avargal/al
6.	Number marker	41, 42, 43, 44	-	-kkal/-gal
7.	Gender marker	45 & 46 47 & 48 49 & 50	Masculine Feminine Neuter	-n -l -u

The categories and markers tested in the both the production task (Picture-live description task and general conversation task)

Data collection

Before commencing the recording, the children were made to feel comfortable. A routine interaction was established with them to achieve rapport so that eliciting responses from them could be smooth. The time taken to record each child was 2 ½ to 3 hours including the rapport building and completion of both the tasks. In the picture description task, the children were instructed to describe the picture in response to the question asked by the investigator. Reinforcements such as verbal and social reinforcements were given at a continuous schedule and tangible reinforcement was given at fixed ratio schedule to encourage participation of the children.

Transcription

The responses of both the tasks for each child in the clinical group and the control group were transcribed verbatim and they were analysed for the presence or absence of the markers/structure in question. The phonological processes and the dialectal variations were not taken into account. To establish the validity of the analysis, the transcripts were reviewed by a trained Linguist.


Means of scoring for the tasks given

When the child gave the expected responses to the picture description task and general conversation it was considered as a correct response and a score of 1 point was given to both the tasks. When the child gave the expected response in one task and not in the other, cues such as binary choice, prompting and semantic cueing was provided. A correct response with the cues was also scored as 1. When the child gave irrelevant answers in both the tasks in spite of the cues given, it was considered as incorrect response and the child was scored 0 in both the tasks. These coded data were statistically analysed by using SPSS (Statistic Program for Social Science) software,

VI. RESULTS

The total score obtained in the two tasks by each child in the control group who had an APGAR score of 8 at the 1st min of birth and 5 children in the clinical group who had an APGAR score of 8 at the 2nd minute after birth had a score of 100. This denotes that these children have acquired all the case markers, tense markers, person/pronoun markers, gender markers and number markers in the language. On the other hand 5 children who had an APGAR score of 8 at the 5th minute after birth had a score of 78 and the remaining 5 children who had an APGAR score of 8 at the 10th minute after birth had a score of 66. This denotes that children who had an APGAR score of 8 at the 5th and 10th minute after birth did not acquire certain markers in the language.

Figure 2




The total score obtained by the children in both the picture-live description and general conversation tasks

The total score obtained by the children in the control and the clinical group in the picture – live description and general conversation tasks. The bar chart indicates that the children who are in the control group (5 children who had an APGAR score of 8 at the 1st min of birth) and 5 children in the clinical group who had an APGAR score of 8 at the 2nd min of birth obtained a score of 50/50 in both picture-live description and general conversation tasks. However 5 children who had an APGAR score of 8 at the 5th min had a score of 39/50 in the picture–live description and general conversation tasks and the remaining 5 children who had an APGAR score of 8 at the 10th min had a score of 33/50 in the picture–live description and general conversation tasks.

Results based on the individual markers present in the language across the tasks

Figure 3



Overall markers attained by the control and clinical group in the production tasks

Figure 3 indicates the total score obtained by the 15 children in the control group (those who had an APGAR score of 8 at the 1st min of birth) and 5 children in the clinical group (those who had an APGAR score of 8 at the 2nd min of birth) for the individual markers in the picture–live description and general conversation tasks were 100. However the total score obtained by the children who had an APGAR score of 8 at the 5th min of birth for the individual markers in both the tasks were 100 for the present tense markers, past tense markers, number markers and gender markers and had obtained a score of 71.4 for case markers, 0 for future tense markers and a score of 70 for person/pronoun marker present in Tamil. The total score obtained by the children who had an APGAR score of 8 at the 10th min of birth for the individual markers in both the tasks were 100 for the present tense markers and number markers and had obtained a score of 71.4 for case markers, a score of 0 for future tense markers, a score of 80 for present tense markers, a score of 70 for person/pronoun markers and a score of 33.3 for gender markers present in Tamil.

Overall Result

Table 7

S.NO	AGE	GENDER	Time at which APGAR score of 8 was obtained	Language Status	
1.	4:0	M	1 st min	All the Case markers, Tense markers, Person/Pronoun markers, Number markers and Gender markers were acquired	
2.	4:1	M	1 st min		
3.	4:2	M	1 st min		
4.	4:2	F	1 st min		
5.	4:4	F	1 st min		
6.	4:7	M	1 st min		
7.	4:8	M	1 st min		
8.	4:8	M	1 st min		
9.	4:9	F	1 st min		
10.	4:11	F	1 st min		
11.	5:0	M	1 st min		
12.	5:4	M	1 st min		
13.	5:6	M	1 st min		
14.	5:3	F	1 st min		
15.	5:5	F	1 st min		
16.	4:0	M	2 nd min		
17.	4:0	M	2 nd min		
18.	4:5	M	2 nd min		
19.	4:3	F	2 nd min		
20.	4:5	F	2 nd min		
21.	4:5	M	5 th min		Except Case marker (-ai), (-ilirundu), Person/Pronoun marker (avar/ar), (avargal) and Future Tense marker (-pp-), (-v-) all the other markers were acquired
22.	4:9	M	5 th min		
23.	4:11	M	5 th min		
24.	4:8	F	5 th min		
25.	4:8	F	5 th min		
26.	5:2	M	10 th min	Except Case marker (-ai), (-ilirundu), Person/Pronoun marker (avar/ar), (avargal), Future Tense marker (-pp-), (-v-), (-in-) and Gender marker (-n), (-l), all the other markers were acquired	
27.	5:5	M	10 th min		
28.	5:6	M	10 th min		
29.	5:1	F	10 th min		
30.	5:6	F	10 th min		

The language status of control and clinical group across both the picture-live description task and the general conversation task

Table 7 indicates the language status obtained from the control and clinical groups with respect to picture description task and general conversation task across their age, gender, time at which APGAR score reached 8.

Table 8

Dependent Variable	Group I - Control group	Group J- Experimental group	Mean difference	Standard error	P value
Total score for Picture-Live Description Task and General Conversation Task	Children with an APGAR score of 8 at the 1 st min of birth	Children with an APGAR score of 8 at the 2 nd min of birth	0.000	0.103	1.000
	Children with an APGAR score of 8 at the 5 th min of birth		0.571*	0.103	0.000
	Children with an APGAR score of 8 at the 10 th min of birth		0.571*	0.103	0.000

Post hoc analyses -Tukey's HSD test to check the language deviance of the children with birth asphyxia by comparing their performance in both the production tasks with age and gender matched control group

Table 8, indicates a high significant value obtained for children whose APGAR score was 8 at the 5th and 10th min when compared to the control group in the picture description task. However there was no significance found in children whose APGAR score was 8 at the 2nd min when compared to the control group in the picture-live description task and general conversation task.

Table 9

Variable	r value	p value
Picture-Live description task	-0.952**	0.000**
General Conversation task	-0.952**	0.000**

** correlation is significant at the 0.01 level (2-tailed)

* correlation is significant at the 0.05 level (2-tailed)

Pearson's correlation to investigate the relation between the time at which the APGAR score reached 8 to the age of the children, performance of the picture description task and general conversation tasks

This table above indicates that when the delay in birth cry increases the performance or the score obtained in both the general conversation task and the picture description task decreases. This shows a negative correlation which is inversely proportional to the variables taken. This finding was highly significant. The pearson's correlation, r value was -0.952** and the p value was 0.000**.

Table 10

Variable	P value
Time at which the APGAR score reached 8	0.010***
Picture-live description task	0.014***
General Conversation task	0.014***

Independent sample test to check the performance of the females both in the control and experimental group across the production tasks

This table indicates that the females had obtained a highly significance value of 0.014** with respect to the 2 tasks.

Table 11

Variable	P value
Time at which the APGAR score reached 8	0.001***
Picture-live description task	0.002***
General Conversation task	0.002***

Independent sample test to check the performance of the males both in the control and experimental group across the production tasks

This table indicates that the males had obtained a highly significance value of 0.002** obtained in males with respect to the 2 tasks.

VII. DISCUSSION

In this study the children who had an APGAR score of 8 at the 1st and 2nd minute exhibited a total score of 100 (maximum score) from both the tasks tested. This denotes that these children had acquired all the case markers (-ai, -kku, -a:l,

-oDa/-kuDa, -ilirundu, -o:da and -il), present tense markers (-kkir-), Future tense markers (-pp- and -v-), Past tense marker (-tt-, -nt-, -d-, -in- and -i-), Person/pronoun marker (na:n, na:ngal, ni:, ni:ngal, a:van, a:val, adu, avar/a:r and avargal/al), number marker (-kkal/-gal), and gender marker (-n, -l and -u). On the other hand the 5 children who had an APGAR score of 8 at the 5th minute exhibited a total score of 78 from both the tasks. This means that these children have not acquired all the evaluated morpho-syntactic units. These children have acquired all except case marker (-ai), (-ilirundu), person/pronoun marker (avar/a:r), (avargal) and future tense marker (-pp-), (-v-). They have acquired all the other markers. Those children (5 children) who had an APGAR score of 8 at the 10th minute exhibited a total score of 66 from both the tasks. This denotes that these children have acquired all the markers in the language except case marker (-ai), (-ilirundu), person/pronoun marker (avar/a:r), (avargal), future tense marker (-pp-), (-v-), past tense marker (-in-) and gender marker (-n), (-l).

When this finding were correlated with the earlier pilot study it was seen that the children who had an APGAR score of 8 at the 5th minute after birth had a language delay of 12 months (Combined Language Age) where as the children who had an APGAR score of 8 at the 10th minute after birth had a language delay of 15 to 24 months (Combined Language Age). This finding could be either because these children were not given any therapeutic intervention to enhance language or it may be because of the delay in the onset of birth cry. In both of these studies, that is the earlier pilot study and the present follow up study, 4 parameters namely the APGAR score, time of birth cry, cognitive abilities pertaining to language and the language status have been given equal importance. Even in the control group were also because even children with mild developmental delay with a developmental quotient less than 90 and slightly delayed social maturity age attend normal kindergarten and primary school. Only when there a deviance from the other peer groups they join in some special school.

Many studies have given importance to the APGAR score and have stated that children who had an APGAR score of 7 had no associated problems but this does not hold good to all the children because many children who had an APGAR score of 7 have associated problems (Majeed et al, 2007). The APGAR score alone cannot give much information on the level or severity of impairment. Hence the APGAR score and time of birth cry along with other parameters such as the cognitive ability should be considered to arrive at a precise conclusion of the presence or absence of associated problems and to check the degree of severity.

In certain studies the APGAR score and the time of birth cry have been given equal importance. Those children with an APGAR score of 7 and had a brief onset of birth cry, that is, within 5 minute after birth had no associated problems such as neurological and cognitive impairment. However in children who had an APGAR score of 7 and had a prolonged onset of birth cry that is after 5 minute of birth had associated problems such as neurological and cognitive impairment (Odd et al, 2008; Paneth, 2010). Jerneck and Herbst (2001) have also revealed that children who had a low APGAR score at the 5th minute after birth are most likely to have mortality neurological impairment, cognitive impairment or both cognitive and neurological impairment. All these manifestations depend on the level of APGAR score obtained by the children at the 5th minute of birth. However, while

documenting the time of birth cry to check the associated problems such as mortality rate, neurological and cognitive problem, the classification of the time of birth cry as brief (within 5 minute after birth) and prolonged (after 5 minutes of birth) is considered to be sufficient, but to check for finer associated problems such as language development, the time of birth cry should be very accurate while documenting. This is because in this present study and the earlier pilot study done to check the language status in children with birth asphyxia it is seen that the onset of birth cry plays a vital role in differentiating who will have adequate language development and who is most likely to have difficulty in acquiring age adequate language development.

Kavitha (2009) found that the children with learning disability and had history of birth asphyxia had difficulty acquiring all the markers in the language when compared to the typically developing children. From the findings it is clear that, children who had birth asphyxia at birth are prone to have delay in the development of language. This has been supported in a study done by Sidhu et al. (2010) which revealed that children who had risk factors such birth asphyxia and other risks exhibited marked delay in the development of language. Stanton-Chapman et al. (2002) stated that one of the early risk factors for language impairment was birth asphyxia. The language development will be significantly delayed when there are more than one associated problem and depending on the severity of the problem (Odd et al, 2010).

Implications of the study

From this study, children who have survived birth asphyxia and had no associated problems may have language delay. This can be attributed to the level of APGAR score, time of birth cry, cognitive level pertaining to the language development and their language status. Most of the earlier studies reported the impact/consequence of asphyxia with reference to either single, two or a maximum of three parameters from APGAR score, Birth cry, cognition and Language but all of the studies had failed to consider all the factors together. This study is the first of its nature where it had addressed the relationship between all the 4 parameters. In most of the studies done on covering the issues on birth asphyxia, the APGAR score at 1st min, 5th min, 10th min, 15th min and 20th min after birth was given much importance. This is because the APGAR score rarely reach 8 and above at the 2nd min of birth. In this study the 2nd min of birth cry is given equal importance as the 1st, 5th and 10th minute of birth cry after birth.

Limitations of the study

Due to the limited sample size non parametric test namely Pearson's correlation was used to obtain qualitative data and the differences could not be noticed in children with an APGAR score of 8 at 1st minute and at the 2nd minute. Percentages were calculated to indicate the markers attained by the children in the production tasks. However, it did not give much information as the scoring allotted in the production tasks were only binary in nature (0 or 1) and the results of the study could not be generalized to all the children who had mild birth asphyxia.

Dialect variation and phonological processes were not taken into account in this study. Although there were equal number of males and females in the control and the clinical group but when grouped in terms of the time at which the APGAR score reached 8 were not proportionate. Hence the

level of significance was much higher in males than in females.

Video recording could not be done for the clinical group due to interrupted electrical facilities and poor room acoustics. The picture cards and the framed questions to elicit general conversation used in this study were not standardized but there were designed to elicit the appropriate morpho syntactic structures. Familiarization of the tasks and cues such as binary choice, prompting and semantic cueing was permitted to elicit appropriate responses from the child. Socio-economic status should have been documented in both the control and the clinical group as the clinical group had low socio economic status whereas the control group had high socio economic status. The children in the control group are studying in a kindergarten or primary school in Chennai where as the clinical groups are studying in government school where "Activity Based Learning" (ABL) is practiced. Although the control group and clinical group were exposed to both English and Tamil, the level of exposure varied depending on the place where they reside, the immediate linguistic environment and their need to communicate in a particular language say Tamil. Biological maternal factors should have been emphasized.

Future directions

These children can be followed up further to check whether the language delay still persists or to check the language status of these children after providing appropriate therapeutic intervention. Studies can be done on other components of language say pragmatics. More studies should in this field to bring reliable results as birth asphyxia is one of the major high risk factor which is of a great concern these days due to its increasing incidence rate. Upcoming studies should consider and implement the limitations of this study.

VIII. CONCLUSION

The study concludes that 15 children in the control group (the typically developing children, who had an APGAR score of 8 at the 1st min after birth) and 5 children from the clinical group who had an APGAR score of 8 at the 2nd minute after birth had acquired all the case markers, tense markers, person/pronoun markers, gender markers and number markers of the language. In contrast, the remaining 10 children from the clinical group (5 children who had an APGAR score of 8 at the 5th minute after birth and 5 children who had an APGAR score of 8 at the 10th minute after birth) did not acquire certain markers of language. This study also shows a strong association between the APGAR score, time of birth cry, the cognitive abilities pertaining to language and the linguistic aspects in terms of acquiring the morpho-syntactic structures of the language. From the study it was seen that there were no significant age and gender differences as the sample size was small.

Acknowledgment

I would like to specially thank my "Jehovah", Parents, Dissertation Guide: Dr. Vaidyanathan, faculty, sibling, relatives, friends, juniors and well-wishers for their dedicated support.

References

- [1] Addy, D. P. (1982). Birth asphyxia. *British Medical Journal*, 284, 1288-1289. doi:10.1136/bmj.284.6325.1288
- [2] Airede, K. I. (1991). Neonatal seizures and a 2-year neurological outcome. *Journal of Tropical Pediatrics*, 37(6), 313-317. doi:10.1093/tropej/37.6.313

- [3] Anderson, V., Catroppa, C., Morse, S., Haritou, F., & Rosenfeld, J. (2005). Functional plasticity or vulnerability following early brain injury? *Pediatrics*, 116, 1374-1382. doi:10.1542/peds.2004-1728
- [4] Anderson, V., Spencer-Smith, M., Leventer, R., Coleman, L., Anderson, P., Williams, J., & Jacobs, R. (2009). Childhood brain insult: Can age at insult help us predict outcome. *Brain*, 132, 45-56. doi:10.1093/brain/awn293
- [5] Apgar, V. A. (1953). A proposal for new method of evaluation of the newborn infant. *Current Research Anesthesia & Analgesia*, 32, 260 - 267.
- [6] Avery, M. E., & Taesch, H. W. (1984). Perinatal factors in handicapping conditions. In Reynolds, C. R., & Fletcher - Janzen, E. (Eds.), *Concise encyclopedia of special education: A reference for the education of children, adolescents, and adults with disability and other exceptional individuals*. New York, NY: John Wiley & sons Inc. Retrieved from http://books.google.co.in/books?id=7rvZT5J3osYC&pg=PA1541&dq=associated+problems+in+high+risk+children++avery,+1984&hl=en&ei=D2ITdiLAcXorQex95jiDg&sa=X&oi=book_result&ct=result&resnum=1&ved=0CC4Q6AEwAA#v=onepage&q=associated%20problems%20in%20high%20risk%20children%20-%20avery%2C%201984&f=false
- [7] Ballantyne, A., Spilkin, A., Hesselink, J., & Trauner, D. (2008). Plasticity in the developing brain: Intellectual, language and academic functions in children with ischaemic perinatal stroke. *Brain*, 131(11):2975-2985. doi:10.1093/brain/awn176
- [8] Beharelle, A. R., Dick, A. S., Josse, G., Solodkin, A., Huttenlocker, P. R., Levine, S. C., & Small, S. L. (2010). Left hemisphere regions are critical for language in the face of early left focal brain injury. *Brain*, 133, 1707 - 1716. doi:10.1093/brain/awq104
- [9] Bhate, S., & Wilkinson, S. (2006). Aetiology of Learning Disability. *Psychiatry*, 5 (9), 298-301. doi:10.1053/j.mppsy.2006.08.001
- [10] Chelvi, K. (2007). Language levels in high risk children: a follow up study. Unpublished Master's dissertation. Sri Ramachandra University, Chennai.
- [11] Cheung, P. Y., & Robertson, C. M. (2000). "Predicting the outcome of term neonates with intrapartum asphyxia." *Acta Paediatrica*, 89(3), 262-264.
- [12] Donn, S. M., Sinha, S. K., & Chiswick, M. L. (2002). *Birth Asphyxia and the Brain: Basic Science and Clinical Implications*, New York, NY: Futura Publishing Company, Inc. Retrieved from http://books.google.co.in/books?id=ygUASdQTBtC&pg=PA132&dq=Birth+Asphyxia+and+the+Brain:+Basic+Science+and+Clinical+Implications+Edited+by+Steven+M.+Donn+et+al.%282002%29&hl=en&ei=m0GIT7dMoSucPyina4M&sa=X&oi=book_result&ct=result&resnum=1&ved=0CCoQ6AEwAA#v=onepage&q&f=false
- [13] D'Souza, S. W., McCartney, E., Nolan, M., & Taylor, I. G. (1981). Hearing, speech and language in survivors of severe perinatal asphyxia. *Archives of disease in childhood*, 56, 245-252. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1627236/pdf/archdisch00770-0015.pdf>
- [14] Ghai, O. P., Gupta, P., & Paul, U. K. (2004). *Newborn Infants*. Essential Pediatrics, New Delhi, India, Interprint publisher.
- [15] Genzel-Boroviczeny, O., Hempelman, J., Zoppelli, L., & Martinez, A. (2010). Predictive value of the 1-min Apgar score for survival at 23-26 weeks gestational age. *Acta paediatrica*, 99(12), 1790-1794. doi:10.1111/j.1651-2227.2010.01937.x
- [16] Golightly, T. H., Raz, S., & Sander, C. J. (2003). Influence of slight to moderate risk for birth hypoxia on acquisition of cognitive and language function in the preterm infant: A cross-sectional comparison with preterm-birth controls. *Neuropsychology*, 17 (1), 3-13. doi:10.1037/0894-4105.17.13
- [17] Gonzalez, F. F., & Miller, S. P. (2006). Does perinatal asphyxia impair cognitive function without cerebral palsy? *Archives Disease in Child Fetal Neonatal*, 91, 454-459. doi:10.1136/adc.2005.092445
- [18] Gustafsson, P., & Kallen, K. (2010). Perinatal, maternal, and fetal characteristics of children diagnosed with attention-deficit-hyperactivity disorder: Results from a population-based study utilizing the Swedish Medical Birth Register. *Developmental Medicine & Child Neurology*, 53(3), 263-268. doi: 10.1111/j.1469-8749.2010.03820.x
- [19] Haider, B. A., & Bhutta, Z. A. (2006). Birth asphyxia in developing countries: Current status and public health implications. *Current problem in pediatric adolescent health care*, 36, 178-188. doi:10.1016/j.cpped.2005.11.002
- [20] Handel, M. V., Swaab, H., D'Vries, L. S., & Jongmans, M. J. (2007). Long-term cognitive and behavioural consequences of neonatal encephalopathy following perinatal asphyxia: A review. *European journal of pediatrics*, 166, 645 - 654. doi:10.1007/s00431-007-0437-8

- [21] Heywood, C., & Canavan, A. (1987). Developmental neuropsychological correlates of language. *Language development and disorders: Clinics in developmental medicine*. London, Mackeith/Blackwell.
- [22] Jansson-Verkasalo, E., Valkama, M., Vainionppa, L., Paakko, E., Ilkko, E., & Lehtihalmes, M. (2004). Language Development in Very Low Birth Weight Preterm Children: A Follow-Up Study, *Folia Phoniatrica Logopedica* 56 (2), 108-119. doi:10.1159/000076062
- [23] Jerneck, K. T., & Herbst, A. (2001). Low 5-minute Apgar score: a population based register study of 1 million term births. *The American college of obstetricians and gynecologists*, 98(1), 65-70. doi:10.1016/S0029-7844(01)01370-9
- [24] Kavitha, C. (2009). Verbs and noun morphology in spoken and written language of children with and without learning disability: a study in Tamil. Unpublished Master's dissertation. Sri Ramachandra University, Chennai.
- [25] Khreisat, W. H., & Hababeh, Z. (2005). Risk factors of birth asphyxia: a study at prince ali ben al-hussein hospital, Jordan. *Pakistan Journal of Medical Science*, 21(1), 30-34. Retrieved from <http://pjms.com.pk/issues/janmar05/pdf/article7.pdf>
- [26] Lee, A. C., Mullany, L. C., Tielsch, J. M., Katz, J., Khatri, S. K., LeClerq, S. C., & Darmstadt, G. L. (2008). Risk factors for neonatal mortality due to birth asphyxia in southern Nepal: A prospective, community-based cohort study. *Pediatrics*, 121(5), 1381-1390. doi:10.1542/peds.2007-1966
- [27] Lee, H. C., Subeh, M., & Gould, J. B. (2010). Low Apgar score and mortality in extremely preterm neonates born in the united states. *Acta paediatrica*, 99(12), 1785-1789. doi:10.1111/j.1651-2227.2010.01935.x
- [28] Lust, B. (2006). *Child language acquisition and growth*, Newyork, NY: Cambridge university press publishers.
- [29] Majeed, R., Menon, Y., Majeed, F., Shaikh, N. P., & Rajar, U. D. (2007). Risk factors of birth asphyxia. *Journal of Ayub Medical College Abbottabad*, 19(3), 67-71. Retrieved from <http://www.ayubmed.edu.pk/JAMC/PAST/19-3/16%20Rehana%20Majeed.pdf>
- [30] Mogford, K. & Bishop, D. (1988). Language development in unexceptional circumstances. In Mogford, k. & Bishop, D. (Eds.). *Language development in exceptional circumstances*, Newyork: NY, Churchill Livingstone, Longmen Group.
- [31] Odd, D. E., Rasmussen, F., Gunnell, D., Lewis, G., & Whitelaw, A. (2008). A cohort study of low Apgar scores and cognitive outcomes. *Archives Disease of Child Fetal Neonatal*, 93, 115 - 120. doi:10.1136/adu.2007.123745
- [32] Odd, D. E., Doyle, P., Gunnell, D., Lewis, G., Whitelaw, A., & Rasmussen, F. (2008). Risk of low Apgar score due to maternal socioeconomic position: A study of Swedish male population. *Acta Paediatrica*, 97 (9), 1275 -1280. doi: 10.1111/j.1651-2227.2008.00862.x
- [33] Odd, D. E., Whitelaw, A., Gunnell, D., & Lewis, G. (2010). The association between birth condition and neuropsychological functioning and educational attainment at school age: a cohort study. *Archives of Disease in Childhood*, 96(1), 30-37. doi 10.1136/adc.2009.176065
- [34] Ohl, C., Czajka, L. D. C., Chobaut, J., & Tavernier, L. (2009). Newborn hearing screening on infants at risk. *International Journal of Pediatric Otorhinolaryngology*, 73, 1691-1695. doi:10.1016/j.ijporl.2009.08.027
- [35] Owens, Jr., R. E. (2001). *Language Development: An Introduction*. (5th ed.), New York, NY: Allyn & Bacon.
- [36] Paneth, N. (2010). Apgar score and risk of cerebral palsy. *British Medical Journal*, 341, 5175. doi:10.1136/bmj.c5175
- [37] Park, A. M., Sanders, T. A., & Maltepe, E. (2010). Hypoxia-inducible factor (HIF) and HIF - stabilizing agents in neonatal care. *Seminars in Fetal & Neonatal Medicine*, 15(4), 196 - 202. doi:10.1016/j.siny.2010.05.006
- [38] Ronald, E. H., Jan, J. E., Hill, A., & Wong, P. K., (1986). Cortical visual impairment following birth asphyxia. *Pediatric Neurology*, 2 (3), 133-137. Retrieved from http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6TBD-4859SCS-22&_user=10&_coverDate=06%2F30%2F1986&_rdoc=1&_fmt=high&_orig=gateway&_origin=gateway&_sort=d&_docanchor=&view=c&_seArchStrId=1685669602&_rerunOrigin=google&_acct=C000050221&_version=1&_urlVersion=0&_userid=10&md5=4a38ad499f3c0ea8fc6c971e225ddcce&searchtype=a
- [39] Shanthini, S. V., & Athmacharan, V. (2009). Language status in children with birth asphyxia: a pilot study. *37th All India Dravidian Linguistic Association, Chennai, India*.
- [40] Shulman, B. B., & Capone, N. C. (2010). Historical and contemporary views of the nature - nurture debates: A continuum of perspectives for the speech - language pathologist. *Language development: foundations, processes and clinical implications*. USA: Jones & Bartlett publishers LLC. Retrieved from http://books.google.co.in/books?id=V3bcMay7wxC&pg=PA72&dq=language+development:+foundations,+processes+and+clinical+implications+%28pp.+61%29.&hl=en&ei=g3yHTbGrL8nMrQecxags&sa=X&oi=book_result&ct=result&resnum=1&ved=0CC0Q6AEwAA#v=onepage&q&f=false
- [41] Sidhu, M., Malhi, P., & Jerath, J. (2010). Multiple risks and early language development. *The Indian journal of pediatrics*, 77(4), 391 - 395. doi:10.1007/s12098-0100044-y
- [42] Simon, E. N., & Morley, G. M (2005). Birth brain injury: Etiology and prevention, Part 111: Concealed and Clandestine trauma, *Medical Veritas*, 2, 513-520. Retrieved from <http://www.autism-end-it-now.org/Downloads/BirthBrainInjuryIII.pdf>
- [43] Somefunu, O. A., Lesib, F. E., Danfulania, M. A., & Olusanya, B. O. (2005). Communication disorders in Nigerian children. *International Journal of Pediatric Otorhinolaryngology*, 70(4), 697-702. doi:10.1016/j.ijporl.2005.09.003
- [44] Stanton-Chapman, T. L., Chapman, D. A., Bainbridge, N. L., & Scott, k. G. (2002). Identification of early risk factors for language impairment. *Research in developmental disabilities*, 23(6), 390-405. doi:10.1016/s0891-4222(02)00141-5
- [45] Straube, S., Voigt, M., Jorch, G., Hailler, E., Briese, V., & Borchardt, U. (2010). Investigation of the association of Apgar score with maternal socio-economic and biological factors: An analysis of German perinatal statistics. *Archives of Gynecology & Obstetrics*. 282, 135-141. doi:10.1007/s00404-009-1217-7
- [46] Taylor, H. G., & Alden, J. (1997). Age related differences in outcomes following childhood brain insults: An introduction and overview. *Journal of International Neuropsychology Society*, 3, 555-567.
- [47] Taylor, H. G., Yeates, K., Wade, S., Drotar, D., Stancin, T., Minich, N. (2002). A prospective study of short and long term outcomes after traumatic brain injury in children : Behaviour and academic achievement. *Neuropsychology*, 16, 15- 27.
- [48] Thomson, A. J., Searle, M., & Russell, G. (1977). Quality of survival after severe birth asphyxia. *Archives of disease in childhood*, 52, 620-626. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1544624/pdf/archdisch00814-0028.pdf>
- [49] Wiig, E. H., & Secord, W. A. (1982). Language disabilities in school age children and youth. In Shames, G. H., Wiig, E. H., & Secord, W. A. (Eds.), *Human communication disorders: An introduction*, MA, Allyn and Bacon.
- [50] Windle, W. (1969). Brain Damage by Asphyxia at Birth. *Scientific American Journal*. 221(4), 76-84.



Sharon Viola Shanthini is an Audiologist and a Speech Language Pathologist. She born and raised in Brunei Darussalam. She completed her Kindergarten till "O" Level in Brunei Darussalam whereas her higher education Plus I, Plus II, Bachelors in Audiology and Speech Language Pathology, Master of Science in Audiology and Speech Language Pathology in India and Hanen Certified Speech Language Pathologist in It Takes Two to Talk (ITTT) in Singapore. She possess a Lifetime Membership of Indian Speech and Hearing Association (ISHA), Overseas Membership of Speech Language and Hearing Association, Singapore (SHAS) since 2012 till 2013, Associate Membership of Malaysian Association of Speech Language and Hearing (MASH): May 2012-May 2014, Certified Member of the Hanen Centre from March 2013 till October 2014 and Member of GSTF (Global Science Technology Forum) since 29/3/13. She has been working in Brunei Darussalam as a Speech Language Pathologist (SLP) in Ace Creative and SLP Consultant in La Vida since 2012.